

# EEG brainmapping of trance states induced by Monochord and Ritual Body Postures in a ritualistic setting

Fachner, Jörg<sup>1</sup> & Rittner, Sabine<sup>2</sup>

<sup>1</sup> University Witten/Herdecke, Chair for Qualitative Research in Medicine

<sup>2</sup> University Clinic of Heidelberg, Dpt. of Medical Psychology in the Center of Psychosocial Medicine

Cite as:

Fachner, Jörg; Rittner, Sabine (2007). EEG brainmapping of trance states induced by Monochord and Ritual Body Postures in a ritualistic setting. In: Isabelle Frohne-Hagemann (Hg.): Receptive Music Therapy - Theory and Practice, p. 189-202. Wiesbaden: Reichert Verlag.

## Abstract

*Two receptive trance-inducing methods (Body-Monochord; Goodman Ritual body postures with fast rattling) were compared to rest. 28 EEG traces of two differently hypnotisable (pHGS scores: male 6.09; female 7.78) subjects were obtained with a mobile brain imager during a group therapy ritual. Emphasis was put on an authentical group setting. Narratives, video protocol, psychometric measures (PCI and 5-D ABZ Test) and amplitude/significance mapping were explored.*

*Both sound-trance inductions caused an increase of Beta-II-percentage, the posture an additional Theta-percentage increase. While body monochord induced frontal desynchronization of the male subject with increase of Beta-II-percentage, indicating ergotropic trance, the female EEG showed a synchronisation with significant ( $p < 0.001$ ) changes in visual and somato-sensoric regions. Her Alpha-changes might indicate change of processing to a trophotropic trance state. Both subjects showed increase of Beta-II-percentage indicating ergotropic trance.*

*During rattling both subjects exhibited frontal-central increases of Theta-percentage. While female subject exhibited more pre-/frontal Theta increases, male subject exhibited more parietal-occipital Theta increases. Significant Changes ( $p < 0.001$ ) on Beta, spectral edge frequencies and Theta-percentage-increases mark this state. Guttman (1990) observed high and low frequency increases in the DC-EEG and called it „paradoxical arousal“. Our results might support his observation.*

## Introduction

Work with trance-inducing sounds has become an intrinsic part of receptive music therapy in practice. Therapeutically intended altered states of consciousness enable an intensification of intrinsic perception, a weakening of psychic barriers of everyday-life consciousness, a more open association and imagination (Aldridge & Fachner, 2006; Rittner & Hess, 1996).

At the beginning of the eighties, some experienced music therapists in Germany perceived a growing interest in the effects of music from outside Europe and started experimenting with trance-inducing effects of monochromatic sounds of monochord, gong, dijeridoo, sound bowls etc., developing individual empirical concepts (Bossinger & Hess, 1993; Hess & Rittner, 1996a; Oelmann, 1993; Rittner & Hess, 1996; Strobel, 1988, 1994; Timmermann, 1983, 1996). These first approaches were revised, further developed and also reviewed critically (Hartogh, 2001; Hess, 1999; Jungaberle, 2003; Rittner & Jungaberle, 2002; Zeuch, 1999). Although the brain plays a decisive role in the experience of

trance states, there is an astonishing lack of music therapy research into psychophysiological aspects of sound-induced trance.

We'd like to present an excerpt from an analysis of two trance inductions, i.e. one in recumbent position on a resounding body monochord (Rittner, 1997) and one "ritual body posture" according to Felicitas Goodman (1989, 1992) accompanied by fast rattling sounds. The whole pilot study analysed four different methods of sound trance in a multi-perspective approach. The research design comprised visual and quantitative evaluations of a spontaneous EEG as well as psychometric measurements ("questionnaire on extraordinary states of consciousness" 5D-ABZ by Dittrich, Lamparter and Maurer, (2002) and "Phenomenology of Consciousness Inventory" (PCI) by Pekala (1991a, 1991b). In addition we did a qualitative, content-analytical evaluation of written reports by participants on their experience.

This article presents the EEG evaluation of two induction methods. An extended electronic journal article (Fachner & Rittner, 2004) containing the coloured EEG brainmaps and audiovisual material can be accessed from the March 2004 issue available at <http://www.musictherapytoday.com>. In German it is available as an online article (Rittner & Fachner, 2004) of the 'Musiktherapeutische Umschau' at <http://www.musiktherapie.de/index.php?id=193>.

## **EEG**

The electroencephalogram (EEG), discovered by Berger in 1929, has lost nothing of its fascination (Berger, 1991). Positron emissions tomography (PET), functional magnet resonance tomography (fMRT) and other procedures produce highly soluble images of the living brain; but the EEG with its temporal exactness is an ideal instrument to measure electrical processes in the brain, especially when listening to music. In addition, an EEG brainmapper provides a map of current intensities and wave speeds. Topographical presentations of main frequency ranges permit conclusions on functional interactions of brain regions and their levels of activity (Maurer, 1989). Slow and large high-amplitude waves (Delta and Theta) e.g. are predominant in sleep, while in states of high concentration and alertness fast waves with small amplitudes (Beta) are dominant. Depending on vigilance and activity, there are various combinations of different waves. The Alpha-frequency band (8-12 Hz), the waves of which emerge when closing one's eyes, is discussed as a mediator and indicator between the high and low frequencies of the EEG, between the vigilant states of waking and sleeping (Basar *et al.*, 1997; Schwendtner-Berlin *et al.*, 1995).

## **Body Monochord**

The body monochord is based on the design of the monochord invented about 15 years ago as a musical instrument for therapy purposes. It has the form of a double-walled wooden stretcher, with 26 strings of equal length and exactly the same tuning on the underside, so that a person lying on the instrument with eyes closed may perceive sound with the entire body, via skin, bones, vibration of

body liquids, and auditory sense. Based on the keynote a, various “sound clouds” with overtones may be produced depending on the type of playing.

### **Ritual body posture with rattle induction**

Primarily, the method of “ritual body posture and ecstatic trance”<sup>®</sup> is not so much a western music therapy procedure but rather an approach from anthropological research providing access to altered states of consciousness, where sound is used as an essential element of trance induction (see also Aldridge & Fachner, 2006; Rouget, 1985). An integration of this method into the clinical practice of music therapy and psychotherapy has turned out to be highly successful (see also Risch *et al.*, 2001; Rittner & Jungaberle, 2002).

A common factor of “ritual body postures” -as they are called by the anthropologist Felicitas Goodman- they are up to 32.000 years old – is that as a rule they are perceived as a considerable effort. In combination with the sounds of a rattle or frame drum of approximately 210 bpm (beats per minute) they produce a spontaneous burst of energy in the body. This allows inducing a controllable hallucinatory experience in the subject on all levels of perception (vision, auditory sense etc.).

The members of our investigation group assumed a specific sitting body posture to which they were not accustomed; Goodman studied this posture entitled “olmecic prince” (see Goodman, 1989: 158-161; Gore, 1996: 170-174). This posture was maintained for 15 minutes, accompanied by the rhythmic high-frequency sound of two calabash rattles, beaten with 210 bpm.

---Insert figure 1---

*Figure 1: Archaeological artefact of the so-called “olmecic prince” (location of find Tabasco/Mexico, 1100-600 B.C. (Source: Gore, 1996: 170).*

---Insert figure 2---

*Figure 2: Test person Tom in the body posture of the “olmecic prince”*

### **Trance and EEG**

The term “trance” (from lat. transire, change over to) has been defined in various, sometimes contradictory ways in literature (Fachner, 2006; Meszaros *et al.*, 2002; Pekala & Kumar, 2000; Rittner, 1999; Rouget, 1985). We have used it here as a generic term for “*various physical-mental alterations that may occur in persons at altered states of consciousness independent of the cultural setting. Stimulus, techniques and ritual that induce and structure a trance depend on the socio-cultural context.*” (Hess & Rittner, 1996b: 395). No EEG analyses on trance experience on the body monochord were found so far. Studies on DC-EEG and ritual body posture were published by Guttman (1990, 1992, 1992).

Available studies on trance and pertinent alterations in the EEG did not focus on music and trance but reviewed individual differences in trance experiences

between a variety of mainly verbal trance inductions (De Benedittis & Sironi, 1985; Jaffe & Toon, 1980; Meszaros *et al.*, 2002; Oohashi *et al.*, 2002; Park *et al.*, 2002; Sabourin *et al.*, 1990). It is interesting to note that more recent studies by Meszaros, Park and Oohashi come closer to the authentic situation, i.e. trance induction in situ, in their research design.

The question whether trance is a temporary state or a specific quality in persons addresses the state-trait discussion in psychology and has not been answered so far (see Meszaros *et al.*, 2002: 500). Most studies assume differences in susceptibility to hypnosis, and subjects were psychometrically differentiated at the beginning. Sabourin e.g. compared 12 persons with high and 12 with low susceptibility to hypnosis respectively. At rest as well as under hypnosis, persons with high susceptibility to hypnosis had higher Theta amplitudes compared to persons with low susceptibility (Sabourin *et al.*, 1990). For an overview and discussion on the above-mentioned studies see Fachner (2006).

### **Aims of the study**

For the purpose of this study we assume that a specific state of consciousness may be induced and that the characteristics emerging in the EEG may be differentiated.

*“In musical psychotherapy with sound trance, music (...) is effective in two directions: 1. Physiologically stirring (ergotropic) towards ecstasy by intensified rhythm in the field of perception (...) or 2. Physically calming and internalizing (trophotropic) towards enstasis with reduced field of perception and focussing via monochromatic sounds”* (Hess & Rittner, 1996b: 401).

Ergotropic trance (*ecstasis*) in this context, in the sense of Fischer’s mapping (Fischer, 1971, 1976; Fischer, 1998), means an alert, non-contemplative, wide-awake altered state of consciousness.

Fischer says: *“The mapping follows along two continua: the perceptive-hallucinatory continuum of increasing central-nervous (ergotropic) excitement, and the perceptive-meditative continuum of increasing (trophotropic) damping.”* (Fischer, 1998). *“Along the two continua, the sensory/motoric ratio increases. This means: the further you go along one continuum, the less will it be possible to verify the sensory element through random motoricity.”* (the same, p. 51)

Accordingly, a trophotropic trance is characterized by a rather relaxed, contemplative, apparently sleepy state, and also by a rather inhibited movement profile, and reduced reaction and willingness to perform (*enstasis*).

This study compares a presumably trophotropic induction (body monochord) with a presumably ergotropic induction (ritual body posture with rattling) in two test persons.

The EEG was expected to show differences of a trophotropic state via EEG synchronization, i.e. via deceleration of the main frequencies and increase of slower wave ranges: Delta from 0.3-4 Hz, Theta from 4-8 Hz and lower Alpha

frequencies of 8-10 Hz (compare David *et al.*, 1983; Schwendtner-Berlin *et al.*, 1995).

Ergotropic states were expected to show in an EEG desynchronization and dominance of high-frequency waves (upper Alpha waves from 10-12 Hz, Beta-I from 12-16 Hz, and Beta II from 16-30 Hz).

Our research questions:

1. Are there intra-individual alterations in the topographical spontaneous EEG compared to undisturbed rest?
2. Which inter-individual differences or common factors may be detected between the two test persons?
3. Does our study reveal an increase in Theta waves in persons highly susceptible to hypnosis?

### **Research Methods**

Topographical alterations of brain activities in two test persons were measured in a group setting with a topographical quantitative EEG brainmapping. Using a pre/post design, we compared artefact-free means of undisturbed rest (baseline state) with sound trance phases (altered state) induced by the body monochord or the ritual body posture with rattling.

#### **Qualitative aspects of set and setting in this electrophysiological study**

It is a well-known difficulty in physiological measurements that movements or activities in the course of a therapy may interfere with the precision of measurement data. EEG brainmapping requires a limitation to receptive sound perception with little movement. Ideally, the test persons were to be recumbent or sitting still in order to avoid movement artefacts in measurement results. The mobile NeuroScience BrainImager® used here was designed for EEG on intensive wards where reliable data collection is essential. Galvanic current separation and filter systems permit the recording of a solid EEG signal.

Measurements were taken in a therapy group setting (N=10). We attached specific importance to measurements being made in the ritual setting of a group well known to the test persons. Unlike an isolated laboratory situation, a group setting and familiarity with the experience will ensure that supportive socio-physiological factors influence the sound-induced trance experience for all participants (naturalistic design).

Most attempts to locate practical music therapy in a laboratory setting impair the authenticity of the situation. The documentation of significant moments in therapy on recording appliances in particular demands a sensitive approach. In the realization of such a qualitative electrophysiology study (see Fachner, 2004) the measuring instruments must be adjusted as close as possible to every-day practice in order to generate explorative data. Consequently, we collect our data in the immediate therapy situation and not under lab conditions. On the basis of such explorative data collection, later a test concept may be designed to review

the tendencies revealed in exploration in a laboratory experiment under ideal technical conditions. Details and discussion may be obtained in Fachner (2001, 2004), Fachner and Rittner (2004) and Burgess & Gruzelier (1997).

### **Test persons**

For this pilot study we selected volunteers with trance experience who were acquainted with the induction methods. The intention was to ensure a high degree of familiarity with the ritual setting and as little sensitivity as possible to the research situation. We focus mainly on the results for two test persons (TP), out of ten participants in total (male TP 1: Tom, female TP 2: Angelika).

### **Study design**

After each of the four trance inductions, the test persons reported their experience in a written journal from which we quote. Two turns were performed, with an interval in between; each time a test person respectively was connected to the mobile brain imager in the group setting. Tom came first, Angelika in the second turn.

We examined a total of four different, tested, receptive music therapy procedures, the trance-inducing effects of which on the participants were measured and compared in the following order:

1. body monochord; designed by H.P. Klein (see Rittner, 1997)
2. singing of monochromatic vocal sounds (see Rittner, 1998)
3. Peruvian Whistling Vessels; old Peruvian whistles, trad. according to Statnekov (2003)
4. ritual body posture with rattle induction (according to Goodman, 1989, 1992).

## **Results**

### **Susceptibility to hypnosis**

The PCI test that quantifies the occurrence of characteristic structures of altered states of consciousness via 12 main dimensions and 14 subscales is a method to determine retrospectively an individual's degree of hypnotic susceptibility (Pekala, 1991a, 1991b). The hypnoidal score indicates in how far the experience of a situation resembles the experience of highly suggestible persons during hypnotic induction. The score for Tom on the integrated "Predicted Harvard Group Scale" (pHGS) shows him as "moderately hypnotizable" with 6,09, while Angelika may be termed "highly hypnotizable" with 7,78 points.

### **Body monochord**

For the rest EEG (baseline) and the subsequent sound EEG (altered state), the test persons were placed with their backs on the body monochord, with a cap on their head containing 28 electrodes, from which cables transferred the data to the brainmapping computer next door.

---insert figure 3---

*Figure 3: Test person with cap lying on the body monochord*

#### *Neuropsychological functions*

Both halves of the brain can be divided into a frontal and a posterior half by the central fissure in the middle. The functions of the two frontal “quarters” may be generally be described as motoric, intentional, planning, and as the hypothetic effective location of the “self”. The two posterior “quarters” have primarily sensory and receptive functions, are in charge of alertness-related processes and visual-spatial orientation. Details on functional correlations of brain regions may be followed up in the pertinent literature, i.e. by Kolb & Whishaw (1996).

#### *EEG alterations TP 1 (Tom)*

Compared to rest, the monochord derivation for Tom showed a decrease of the frontal Delta waves and parietal Theta and Alpha and Beta II waves. However, the fast Beta-II frequencies (16-30Hz) showed an increase in the frontal regions. Accordingly, the t-test comparison between rest and monochord revealed highly significant ( $p < 0.001$ ) differences on the Beta II frequency band from frontal to Gyrus Praecentralis.

The spectral band presenting the dominant peak frequencies and changes in frequency speeds showed an advancing desynchronization (increase of frequency speed) compared to rest.

While Theta and low Alpha frequencies were observed in frontal regions at rest, the monochord phase was characterized by the onset of a desynchronization, with a dominance of medium and high Alpha frequencies. This change in frontal frequency speeds was highly significant.

#### **EEG changes TP 2 (Angelika)**

In the case of Angelika, a decrease of Theta waves in temporal and frontal regions occurred compared to rest, while Alpha and Beta frequencies increased in parietal regions. In contrast to Tom a synchronization of the EEG to the frontal regions was recognizable in the spectral band during the monochord phase.

Changes were generally significant on the Alpha band, however not in frontal regions on the Theta band. Moreover, highly significant changes occurred in occipital regions on both Beta bands.

#### **Ritual body posture with rattle induction**

##### *EEG changes TP 1 (Tom)*

With the exception of the amplitude increase on the Beta II and the Alpha band, the amplitude mapping for Tom shows an increase of Theta and Beta I as well as a left-hemispheric and occipital desynchronization in the spectral band. The significance mapping for Tom also resembles that for Angelika.

*EEG changes TP 2 (Angelika)*

Compared to rest, the ritual body posture in Angelika combined with rattling reveals an increase of Theta waves from prefrontal and frontal and parietal-occipital along the vertex. Moreover, Alpha increases are to be found on the right, temporal-occipital and parietal. The Beta bands show distinct increases in parietal-occipital regions. The spectral band also represents a clearly visible increase in left-hemispheric frequency speeds.

In significance mapping, the contrast to rest is represented in highly significant alterations on the spectral, Beta I + II and the Alpha band.

## **Discussion**

### **EEG correlate of the monochord experience**

Both test persons showed an increase in Beta waves and a decrease in Theta waves during the monochord phase.

Diminutions in Theta waves, specifically in temporal regions (where primary auditory centres are located), while subjects were listening to music, were also found in a further study using this topographical EEG method. Fachner (2002) described this effect in a comparison of rest and music. Measurements with a direct-voltage EEG also revealed decreases in temporal regions (Altenmüller & Beisteiner, 1996; David *et al.*, 1969). Fachner found in his study with the NeuroScience BrainImager® that, in addition, the activity in the temporal regions is further dampened after consumption of a psychoactive substance (Cannabis). While the EEG signature of the substance was recognizable in synchronisation and Alpha wave increase, this study appears to show that reaction to trance induction – in the case of the monochord in particular – is rather specific to the individual test person.

Despite growing physical relaxation, the desynchronisation and increase of high Beta frequencies compared to rest suggests an active visual imagination in the case of Tom (*“A cloud rises in me and through the back of my head pulls me up to great heights ... I am flying, surrounded by clouds ... thousands of houses in green and lilac on a slope ... a beautiful sight...”*). The simultaneous decrease of Theta waves suggests increasing physical-mental alertness and an intentionally controlled imagination. The test person Tom influences his rather ecstatic experience.

In contrast, Angelika’s EEG tends towards synchronization, induced by the Alpha increases. For her, the recumbent position on the monochord appears to promote a more relaxed and sensual body experience (*“.. in me this turning movement .. as if I hovered in this rotation ... there were patches of haze or drifting veils and far away an unearthly music ... a feeling of calmness, of being sheltered...”*). She seems to “drift” into an enstatic experience, a contemplation and deep physical relaxation. This was most obvious in the t-test in the highly significant change of the Alpha frequency in the comparison of rest and trance. Accordingly, the trance phase EEG revealed an increase in Alpha waves. Crawford underlined the correlation between Theta activity, high suggestibility

and a reaction of frontal and limbic regions. Function-related hippocampus and amygdala activities (in deeper layers of the central brain) suggest a complex pattern of facilitation and inhibition of neural interaction in the limbic system of persons highly suggestible to hypnosis (in Sabourin *et al.*, 1990). Since our two subjects turned out to differ in their suggestibility to hypnosis as a test result, i.e. revealed differences in facilitation and inhibition, this seems to influence the experience regarding tendencies towards ecstatic or enstatic experience. For Angelika, the hypnoidal score of the pHGS indicated “highly hypnotizable” (see above); she reacts more in the sense of a trophotropic trance with increasing low-frequency waves, while Tom reacts in the sense of an ergotropic trance with increasing high-frequency waves.

The personal written reports reflect the effects of the monochord on both test persons in similar, subjective descriptions of hovering states, visions of cloud-like forms and changes in the body feeling. *“My impression is that I drift through space lying on the monochord”* is Tom’s description. Angelika: *“They were very slow, but even movements ... as if I hovered in this rotating movement ... a tremendous space in grey and white, through which veils were drifting”*.

### **EEG correlates of ritual body posture with rattling**

#### *Reports on experience*

How to describe the experience of a “ritual body posture with rattle induction”? Exemplary excerpts from written reports by Tom and Angelika serve as illustrations. Angelika saw and perceived herself as a “vulcano” in trance:

*I see the streams of lava in orange-red seeping over the black earth. It is still happening extremely slowly but evenly. Then I can feel a large opening in my lower abdomen through which lava comes out of me, too. I see it seeping out of me, can feel it in a very sensual way, and at the same time I am lava myself. After a long interval I hear the rattles again. They sound like thundering stones. My hands become rocks, which I throw away. Rocks keep growing, and I keep hurtling them away. I am still the vulcano. All this does not feel very alarming but powerful. Then the movements of hurtling stones and of streaming lava cease. I feel light emerging in front of me. Everything is perfectly still, I am immobile, too. I can see tiny green plants sprouting from my hands, can feel them coming out between my fingers. – I feel full of power and content.”* (Further details in: (Rittner, 2004)

Tom identifies with the archaeological artefact of the “olmecic prince”:

*“As soon as the rattling starts, I start to drift. I see figures and objects in front of me. An altar made of stone, in the middle of jungle. The images are more what I feel than what I see. Something bright and yellow on the altar .. the brightness becomes a crown, like the head dress of the*

*“olmecic prince”. The crown becomes larger and brighter. I see the very tall olmecic figure sitting in front of the altar. Creepers climb up all over it ... the whole place is ... grown over with plants. A tall, upright serpent ... looks at me, in the foreground. This is a very powerful, “knowing” place ... I see the earth as a rotating globe in green and blue. I watch it for some time and then I think: “and this is why the earth moves!” – The rattling stops, far too soon, a pity. I feel “dazed” for a long time. I feel I have taken in a lot of knowledge... the key to essential questions...”*

Experience gained in altered states of consciousness becomes apparent e.g. in changes in thinking, emotionality, perception of meaning and a feeling of something unutterable (Ludwig, 1966). A known effect is improved memory of what has been perceived during the ritual body posture. The increase in Theta waves and highly significant changes on fast frequency bands observed in Angelika, in visual regions specifically, may represent a more active access to image-like structures, analogous to a dream experience. The images and insights perceived or remembered in trance through spontaneous emotional intuition (*“I still am the volcano. All this does not feel very alarming but powerful...”*) are possibly reflected here in the increase of Theta waves in the limbic system and of Beta waves in visual regions.

In 1990, Guttman was the first to demonstrate in studies with the direct-voltage EEG that in the ritual body posture with rattle induction and simultaneously with the occurrence of Theta waves in the EEG, there was an unusual increase of cortical negativation (DC potential) of 2000-3000 microvolt (Guttman, 1990: 319). This indicates an overactive state of the cerebral cortex and in spontaneous EEG would be evident in a dominance of Beta waves. Guttman coined the term “paradoxical arousal” or “relaxed high tension” for this phenomenon.

Park et al. (2002) found changes in the EEG in the case of a Salpuri dancer in comparing rest, listening to music and memory (of a previous dance). Salpuri is a traditional dance performed by medicine men in Korea. In the phase of remembering an ecstatic trance state in the dance, frontal increases of low alpha frequencies (8-10Hz) were found as well as frontal-occipital Theta increases, compared to rest. In the comparison of rest and listening to a piece of pop music, there was a highly significant increase in the frequency of high Alpha frequencies (10-12.5 Hz) over the entire cortex. This may have indicated a difference between a primarily physical trance experience and the enjoyment felt in listening to music. 9.5 Hz was the identified peak frequency in rest and memory of dance, with an increase of amplitude energy in the memory of dance. The peak frequency rose to 10 Hz in the process of listening to music, and high Beta frequencies increased. Park supposes that the Salpuri dancer *“reaches the altered state of ecstatic trance through suppression of frontal cortex functions and activation of subcortical functions”* (Park et al., 2002: 961), i.e. that trance

is characterized by Theta frequencies, the dominance of which in the EEG suggests such activation.

### **Beta changes**

The increase in high Beta frequencies, while a subject listened to music reported by Park is also known from other studies on music perception. Walker (1977) reported increased right-lateral activity while listening to classical music; Behne et al (1988) reported occipital increases; Petsche (1993) found increases in posterior right-lateral coherence, and Bruggenwerth reported music-related, emotion-specific decreases or increases of posterior Beta activity (Bruggenwerth *et al.*, 1994). According to Petsche (1994), the Beta frequency bands indicate differentiations of music-related cognitive activity.

The conspicuous reaction of the EEG in both trance inductions with an increase of high Beta waves (16-30Hz) seems to be a further indication of the influence of trance triggered by sounds. Beta frequencies indicate changes in emotional states. Isotani et al. (2001) explored hypnosis-induced states of relaxation and anxiety and discussed the conspicuous reactions with high Beta frequencies and their EEG signature in emotional states.

Meszaros et al. (2002) interpreted the EEG for hemispheric differences and described a primarily right-hemispheric, parieto-temporal EEG reaction of the Alpha and Beta band in persons highly susceptible to hypnosis; he concluded that in the “*mainly emotion-focussed hypnotherapies*” (the same, p. 511), as expected, right-hemispheric changes will be dominant. In his study, persons with high, medium and low susceptibility to hypnosis all experienced an altered state of consciousness while listening to music in a relaxed position in an easy chair. Significant differences in the “Altered State Index” were only found for the areas imagination/hallucination (compare the same, p. 510).

Consequently, the Beta changes we found for the monochord as well as for the ritual body posture seem to illustrate the emotional reactions produced by the ritual and the sounds, which in the case of Tom resulted in an anterior right-hemispheric, and for Angelika in a posterior left-hemispheric change. It has to be mentioned that the high muscle tension, which is part of the ritual body postures, especially here in the neck region, has surely an influence on beta frequency range. This can be monitored with an EMG as well and should necessarily be investigated in future research. Therefore the interpretation of the Beta range in occipital regions has to be regarded as a tendency.

### **Conclusions**

Returning to the research questions stated at the beginning, we found inter- and intra individual differences in comparing rest and trance induction. In this study with two test persons, highly significant differences were found in the topographical EEG comparing rest and trance. The profile becomes irregular in the case of the monochord. The female subject recumbent on the monochord showed a more trophotropic trance, indicated by an increase in Alpha waves,

while the male subject showed a desynchronization and an increase of Beta waves, which suggests a more ergotropic trance. The data available do not explain whether these findings suggest gender-specific or mood-dependent differences, or differences in susceptibility to hypnosis.

Moreover, the topographical EEG showed a distinct difference between rest and sound-induced activations. The temporal regions of both test persons revealed a decrease on the Theta band.

Both subjects, however, showed similar reactions in trance induction through ritual body posture accompanied by rattling sounds. In both cases, there was a simultaneous increase in low and high frequency waves of the Theta and the Beta band. This state, described by Guttman as *paradoxical arousal*, appears to be a specific reaction to this method of trance induction rediscovered by Goodman. We confirmed for a ritual group setting what Guttman et al. discovered in laboratory tests as early as 1990: an ecstatic trance induced by sound and body posture is characterized by a “relaxed high tension”.

A possible answer to our question concerning Theta increases in easily hypnotizable persons was only visible in body posture; but in the monochord test we also found a tendency toward synchronization and increased low frequencies in the easily hypnotizable subject. Increases in Theta and Alpha waves during the trance phase reported by other authors were most pronounced in the ritual body posture.

These results represent only a very small part of the entire data evaluated and are of an explorative nature. The last measurement phase covers, as expected, influences of the three previous trance inductions via singing, monochord, and listening to and playing of old Peruvian whistles. An ergotropic trance may well be characterized by an increase in high-frequency waves, and a trophotropic trance by an increase of low-frequency waves. The ritual body posture with rattles seems to cover both forms of reactions. To show more than a tendency of the beta range in the posterior regions an EMG has to be recorded during the postures as well.

Contrary to the concern of suffering loss of control in altered states of consciousness, our repeated findings of high Beta II elements illustrate that self-control is maintained in these methods of sound-induced trance and may even give way to a highly concentrated hyper alertness in the sense of an ergotropic trance. This is confirmed by the participating test persons as well as by our test results. A welcome effect of our study would be to reduce the fears deeply rooted in our society for cultural, historical and political reasons that trance is equivalent to loss of control and susceptibility to manipulation (Rittner, 1999).

### **Acknowledgement:**

We thank Prof. Eduard David, M.D., director of the Centre of Electropathology, University Witten/Herdecke, and former Chair of the Institute of Physiology, University Witten/Herdecke, who made the NeuroScience BrainImager® available to us for the purpose of this study.

## References

- Aldridge, D., & Fachner, J. (Eds.). (2006). *Music and altered states - Consciousness, transcendence, therapy and addictions*. London: Jessica Kingsley.
- Altenmüller, E., & Beisteiner, R. (1996). Musiker hören Musik: Großhirnaktivierungsmuster bei der Verarbeitung rhythmischer und melodischer Strukturen. In *Musikpsychologie - Jahrbuch der Deutschen Gesellschaft für Musikpsychologie* (pp. 89-109). Wilhelmshaven: Florian Noetzel.
- Basar, E., Schurmann, M., BasarEroglu, C., & Karakas, S. (1997). Alpha oscillations in brain functioning: An integrative theory. *Int J Psychophysiol*, 26(1-3), 5-29.
- Behne, K. E., Lehmkuhl, P., & Hassebrauck, M. (1988). EEG-Korrelate des Musikerlebens, Teil II. In Behne, K. E., Kleinen, G. & de la Motte-Haber, H. (Eds.), *Musikpsychologie* (pp. 95-106). Wilhelmshaven: Florian Nötzel.
- Berger, H. (1991). *Das Elektroenkephalogramm des Menschen - Kommentierter Reprint des Erstdruckes aus dem Jahre 1938 (Kommentierter Reprint des Erstdruckes von 1938, Halle ed.)*. Frankfurt am Main: PMI Verlag.
- Bossinger, W., & Hess, P. (1993). Musik und außergewöhnliche Bewusstseinszustände. *Musiktherapeutische Umschau*, 14(3), 239-254.
- Bruggenwerth, G., Gutjahr, L., Kulka, T., & Machleidt, W. (1994). Music induced emotional EEG reactions. *EEG-EMG-Z Elektroenz Elektrom*, 25(2), 117-125.
- Burgess, A. P., & Gruzelier, J. (1997). How reproducible is the topographical distribution of EEG amplitude? *Int J Psychophysiol*, 26(1-3), 113-119.
- David, E., Berlin, J., & Klement, W. (1983). Physiologie des Musikerlebens und seine Beziehung zur trophotropen Umschaltung im Organismus. In Spintge, R. & Droh, R. (Eds.), *Musik in der Medizin - Neurophysiologische Grundlagen, Klinische Applikationen, Geisteswissenschaftliche Einordnung* (pp. 33-48). Berlin: Springer.
- David, E., Finkenzeller, P., Kallert, S., & Keidel, W. D. (1969). Akustischen Reizen zugeordnete Gleichspannungsänderungen am intakten Schädel des Menschen. *Pflügers Archiv*, 309, 362-367.
- De Benedittis, G., & Sironi, V. A. (1985). Deep cerebral electrical activity during hypnotic state in man: Neurophysiologic considerations on hypnosis. *Rivista di Neurologia*, 55(1), 1-16.
- Dittrich, A., Lamparter, D., & Maurer, M. (2002). *5D-ABZ - Fragebogen zur erfassung aussergewöhnlicher Bewusstseinszustände (2. ed.)*. Zürich: PSIN PLUS.
- Fachner, J. (2001). *Veränderte Musikwahrnehmung durch Tetra-Hydro-Cannabinol im Hirnstrombild (E-Book on Music Therapy Info CD-ROM ed. Vol. III)*. Witten: University Witten/Herdecke.
- Fachner, J. (2002). Topographic EEG changes accompanying cannabis-induced alteration of music perception - Cannabis as a hearing aid? *Journal of Cannabis Therapeutics*, 2(2), 3-36.
- Fachner, J. (2004). Cannabis, brain physiology, changes in states of consciousness and music perception. In Aldridge, D. (Ed.), *Case Study Designs in Music Therapy* (pp. 211-233). London: Jessica Kingsley.
- Fachner, J. (2006). Music and altered states of consciousness - an overview. In Aldridge, D. & Fachner, J. (Eds.), *Music and Altered States - Consciousness, Transcendence, Therapy and Addictions* (pp. 15-37). London: Jessica Kingsley.
- Fachner, J., & Rittner, S. (2004). Sound and trance in a ritualistic setting visualised with EEG Brainmapping. *Music Therapy Today Online* Retrieved 18. January, 2006, from <http://www.musictherapyworld.net/modules/mmmagazine/showarticle.php?articletoshow=93>

- Fischer, R. (1971). A Cartography of the Ecstatic and Meditative States. *Science*, 174, 897-904.
- Fischer, R. (1976). Transformations of consciousness. A cartography, II. The perception-meditation continuum. *Confin Psychiatr*, 19(1), 1-23.
- Fischer, R. (1998). Über die Vielfalt von Wissen und Sein im Bewusstsein. Eine Kartographie außergewöhnlicher Bewusstseinszustände. In Verres, R., Leuner, H. C. & Dittrich, A. (Eds.), *Welten des Bewußtseins* (pp. 43-70). Berlin: VWB - Verlag für Wissenschaft und Bildung.
- Goodman, F. (1989). *Wo die Geister auf den Winden reiten. Trancereisen und ekstatische Erlebnisse*. Freiburg: Bauer.
- Goodman, F. (1992). *Trance - Der uralte Weg zu religiösem Erleben. Rituelle Körperhaltungen und ekstatische Erlebnisse*. Gütersloh: GTB.
- Gore, B. (1996). *Ekstatische Körperhaltungen*. Essen: Synthesis.
- Guttman, G. (1990). Zur Psychophysiologie der Bewußtseinssteuerung. *Meditation – Trance – Hypnose: Wurzeln und biologische Korrelate*. In Bandion, J. (Ed.), *Einheit der Vielfalt*. Wien: Gerold.
- Guttman, G. (1992). Zur Psychophysiologie des Bewußtseins. In Guttman, G. & Langer (Eds.), *Das Bewußtsein. Multidimensionale Entwürfe*. Wien: Springer.
- Guttman, G., & Langer, G. (1992). *Das Bewußtsein - Multidimensionale Entwürfe*. Wien: Springer.
- Hartogh, T. (2001). Die Rezeption monotonaler Klänge. Eine empirische Untersuchung zur Klangwirkung des Monochords. *Zeitschrift für Musik-, Tanz- und Kunsttherapie*, 12(3), 111-119.
- Hess, P. (1999). Musiktherapie mit archaischen Klangkörpern. *Musiktherapeutische Umschau*, 20(2).
- Hess, P., & Rittner, S. (1996a). Trance. In Decker-Voigt, H. H., Knill, P. & Weymann, E. (Eds.), *Lexikon Musiktherapie* (pp. 395-398). Göttingen: Hogrefe.
- Hess, P., & Rittner, S. (1996b). Verändertes Wachbewusstsein. In Decker-Voigt, H. H., Knill, P. & Weymann, E. (Eds.), *Lexikon Musiktherapie* (pp. 398-403). Göttingen: Hogrefe.
- Jaffe, J. R., & Toon, J. H. (1980). EEG and polygraphic changes during hypnotic suggestibility. *Journal of Electrophysiological Technology*, 6(2), 75-92.
- Jungaberle, H. (2003). Töne strömen ein. Beiträge zur Musikpsychologie des Monochords. *Musiktherapeutische Umschau*, 24(3).
- Kolb, B., & Wishaw, I. Q. (1996). *Neuropsychologie*. Heidelberg: Spektrum Verlag.
- Ludwig, A. M. (1966). Altered States of Consciousness. *Arch Gen Psychiatry*, 15(3), 225 - 234.
- Maurer, K. (Ed.). (1989). *Topographic mapping of EEG and evoked potentials*. Berlin: Springer.
- Meszaros, I., Szabo, C., & Csako, R. I. (2002). Hypnotic susceptibility and alterations in subjective experiences. *Acta Biol Hung*, 53(4), 499-514.
- Oelmann, J. (1993). Klang, Wahrnehmung, Wirkung. Zur therapeutischen Arbeit mit Gongs und Tam-Tams in rezeptiver Therapie. *Musiktherapeutische Umschau*, 14(4), 289-305.
- Oohashi, T., Kawai, N., Honda, M., Nakamura, S., Morimoto, M., Nishina, E., et al. (2002). Electroencephalographic measurement of possession trance in the field. *Clinical Neurophysiology*, 113(3), 435-445.
- Park, J. R., Yagy, T., Saito, N., Kinoshita, T., & Hirai, T. (2002). Dynamics of brain electric field during recall of Salpuri dance performance. *Percept Mot Skills*, 95(3 Pt 1), 955-962.
- Pekala, R. J. (1991a). *Phenomenology of Consciousness Inventory (PCI) Form I ((1999), U. O., Trans.)*. West Chester PA: MID-Atlantic Educational Institute.

- Pekala, R. J. (1991b). *Quantifying Consciousness: an empirical approach*. New York, NY: Plenum Press.
- Pekala, R. J., & Kumar, V. K. (2000). Operationalizing 'trance' I: Rationale and research using a psychophenomenological approach. *American Journal of Clinical Hypnosis*, 43(2), 107-135.
- Petsche, H. (1993). Zerebrale Verarbeitung. In Bruhn, H., Oerter, R. & Rösing, H. (Eds.), *Musipsychologie* (pp. 630-638). Reinbeck bei Hamburg: Rowohlt.
- Petsche, H. (1994). The EEG while listening to music. *Eeg-Emg-Z Elektroenz Elektrom*, 25(2), 130-137.
- Risch, M., Scherg, H., & Verres, R. (2001). Music therapy for chronic headaches. Evaluation of music therapeutic groups for patients suffering from chronic headaches. *Schmerz*, 15(2), 116-125.
- Rittner, S. (1997). Die Arbeit mit dem Ganzkörper-Monochord in der Musikpsychotherapie. In Berger, L. (Ed.), *Musik, Magie & Medizin* (pp. 110-117). Paderborn: Junfermann.
- Rittner, S. (1998). Singen und Trance – Die Stimme als Medium zur Induktion veränderter Bewußteinszustände. In Gundermann, H. (Ed.), *Die Ausdruckswelt der Stimme* (pp. 317 -325). Heidelberg: Hüthig.
- Rittner, S. (1999). Bewußteinsforschung. In Verres, R., Schweizer, J., Jonasch, K. & Süßdorf, B. (Eds.), *Heidelberger Lesebuch Medizinische Psychologie*. Heidelberg: Vandenhoeck & Rupprecht.
- Rittner, S. (2004). Seelen-Wege: Der Körper als Gefährt für Reisen in andere Wirklichkeiten. In Tucek, G. (Ed.), *Musik und Medizin - Beiträge zur Musik- und Therapieforschung 1995 - 2004 (CD-ROM Edition)*. Wien: Akademie für Ganzheitsmedizin.
- Rittner, S., & Fachner, J. (2004). Klang und Trance im EEG – Brainmapping verschiedener Tranceinduktionsmethode im rituellen Setting. *Musiktherapeutische Umschau Online* Retrieved 18. Januar, 2006, from <http://www.musiktherapie.de/index.php?id=193>
- Rittner, S., & Hess, P. (1996). Klangtrance. Trance. Veränderte Bewußteinszustände. In Decker-Voigt, H. H., Weymann, E. & Knill, P. (Eds.), *Lexikon Musiktherapie* (pp. 169-173). Göttingen: Hogrefe.
- Rittner, S., & Jungaberle, H. (2002). Sounding systems - Different Approaches to the Evaluation of Music Psychotherapy Groups. In Aldridge, D. & Fachner, J. (Eds.), *Music Therapy in Europe - Proceedings of the the 5th European Music Therapy Congress in Castel Dell'Ovo, Naples, Italy 20.-24. April 2001 (E-Book on Music Therapy Info CD-ROM ed., Vol. IV, pp. 1514-1543)*. Witten: University Witten/Herdecke.
- Rouget, G. (1985). *Music and Trance. A Theory of the Relations between Music and Possession*. Chicago: University Press.
- Sabourin, M. E., Cutcomb, S. D., Crawford, H. J., & Pribram, K. (1990). EEG correlates of hypnotic susceptibility and hypnotic trance: Spectral analysis and coherence. *International Journal of Psychophysiology*, 10(2), 125-142.
- Schwendtner-Berlin, H., Berlin, J., David, L., Fachner, J., Böving, S. F., David, E., et al. (1995). Gesetzmäßigkeiten in der Psychologie veränderter Bewußteinszustände. *Curare*, 18(2), 361-368.
- Statnekov, D. K. (2003). *Animated Earth (2 ed.)*. Berkeley, CA: North Atlantic Books.
- Strobel, W. (1988). Klang - Trance - Heilung. Die archetypische Welt der Klänge in der Psychotherapie. *Musiktherapeutische Umschau*, 9, 119-139.
- Strobel, W. (1994). Die klanggeleitete Trance. In *Welten des Bewußtseins - Bedeutung für die Psychotherapie (Vol. 4, pp. 225-237)*. Berlin: VWB.
- Timmermann, T. (1983). *Klangstrukturen und ihre psychische Wirkung*. München: Freies Musikzentrum e.V.

- Timmermann, T. (1996). Ethnologische Aspekte in der Musiktherapie. In Decker-Voigt, H. H., Knill, P. & Weymann, E. (Eds.), *Lexikon Musiktherapie* (pp. 87-90). Göttingen: Hogrefe.
- Walker, J. L. (1977). Subjective reactions to music and brainwave rhythms. *Physiological Psychology*, 5(4), 483-489.
- Zeuch, A. (1999). Schöpfer eigener Wirklichkeiten. Die Klangmeditation mit dem Monochord. *Musik-, Tanz- und Kunsttherapie*, 10(4), 175-185.